



Departement of Computer Science

26 September 2022

Markus Püschel, David Steurer

François Hublet, Goran Zuzic, Tommaso d'Orsi, Jingqiu Ding

Algorithms & Data Structures

Exercise sheet 1

HS 22

The solutions for this sheet are submitted at the beginning of the exercise class on 3 October 2022.

Exercises that are marked by * are “challenge exercises”. They do not count towards bonus points.

You can use results from previous parts without solving those parts.

Exercise 1.1 *Guess the formula (1 point).*

Consider the recursive formula defined by $a_1 = 1$ and $a_{n+1} = 2a_n + 1$. Find a simple closed formula for a_n and prove that a_n follows it using induction.

Hint: Write out the first few terms. How fast does the sequence grow?

Exercise 1.2 *Sum of Squares.*

Prove by mathematical induction that for every positive integer n ,

$$1^2 + 2^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}.$$

Exercise 1.3 *Sums of powers of integers (1 point).*

In this exercise, we fix an integer $k \in \mathbb{N}_0$.

- (a) Show that, for all $n \in \mathbb{N}_0$, we have $\sum_{i=1}^n i^k \leq n^{k+1}$.
- (b) Show that for all $n \in \mathbb{N}_0$, we have $\sum_{i=1}^n i^k \geq \frac{1}{2^{k+1}} \cdot n^{k+1}$.

Hint: Consider the second half of the sum, i.e., $\sum_{i=\lceil \frac{n}{2} \rceil}^n i^k$. How many terms are there in this sum? How small can they be?

Together, these two inequalities show that $C_1 \cdot n^{k+1} \leq \sum_{i=1}^n i^k \leq C_2 \cdot n^{k+1}$, where $C_1 = \frac{1}{2^{k+1}}$ and $C_2 = 1$ are two constants independent of n . Hence, when n is large, $\sum_{i=1}^n i^k$ behaves “almost like n^{k+1} ” up to a constant factor.

Exercise 1.4 *Asymptotic growth (1 point).*

Recall the concept of asymptotic growth that we introduced in Exercise sheet 0: If $f, g : \mathbb{N} \rightarrow \mathbb{R}^+$ are two functions, then:

- We say that f grows asymptotically slower than g if $\lim_{m \rightarrow \infty} \frac{f(m)}{g(m)} = 0$. If this is the case, we also say that g grows asymptotically faster than f .

Prove or disprove each of the following statements.

(a) $f(m) = 100m^3 + 10m^2 + m$ grows asymptotically slower than $g(m) = 0.001 \cdot m^5$.

(b) $f(m) = \log(m^3)$ grows asymptotically slower than $g(m) = (\log m)^3$.

(c) $f(m) = e^{2m}$ grows asymptotically slower than $g(m) = 2^{3m}$.

Hint: Recall that for all $n, m \in \mathbb{N}$, we have $n^m = e^{m \ln n}$.

(d) $f(m) = \sum_{i=1}^{m^2} i$ grows asymptotically slower than $g(m) = \sum_{i=1}^m i^2$.

Hint: You can reuse the inequalities from exercise 1.2.

(e)* If $f(m)$ grows asymptotically slower than $g(m)$, then $\log(f(m))$ grows asymptotically slower than $\log(g(m))$.

(f)* $f(m) = \log(\sqrt{\log(m)})$ grows asymptotically slower than $g(m) = \sqrt{\log(\sqrt{m})}$.

Hint: You can use L'Hôpital's rule from sheet 0.

Exercise 1.5 Proving Inequalities.

(a) By induction, prove the inequality

$$\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdots \frac{2n-1}{2n} \leq \frac{1}{\sqrt{3n+1}}, \quad n \geq 1.$$

(b)* Replace $3n + 1$ by $3n$ on the right side, and try to prove the new inequality by induction. This inequality is even weaker, hence it must be true. However, the induction proof fails. Try to explain to yourself how is this possible?